

Effectiveness of Rat Bone Marrow Stem Cell Therapy to *Rattus novergicus* by Teratogenic Model of Particulate Matter on Expression of Kappa Beta (NFk(I) Nuclear Factor on Placenta

by Bu Rima

Submission date: 30-Sep-2019 10:46AM (UTC+0800)

Submission ID: 1182692236

File name: 16.Effectiveness_of_Rat....pdf (1.28M)

Word count: 786

Character count: 4477

Effectiveness of Rat Bone Marrow Stem Cell Therapy to *Rattus novvergicus* by Teratogenic Model of Particulate Matter on Expression of Kappa Beta (NF κ B) Nuclear Factor on Placenta

Sri Pantja Madyawati^{1*}, Widjiati², Rimayanti³

¹Laboratory of Reproduction Physiology, Faculty of Veterinary Medicine, Airlangga University

²Laboratory of Embryology, Faculty of Veterinary Medicine, Airlangga University

³Laboratory of Infertility, Faculty of Veterinary Medicine, Airlangga University

*Corresponding author: sripantja_madyawati@yahoo.com

Key words: expression of NF- κ B, Rat Bone Marrow Stem Cell, teratogenic, particulate matter

INTRODUCTION

Air pollution in several cities in East Java has exceeded the threshold set by the government. One of the pollutant components that pollute the air is particulate matter (PM) [1]. PM component effects health because it is cytotoxic and teratogenic. Particulate Matter can cause inflammation on the respiratory and cardiovascular systems, and it can pass through the placental barrier so that it can affect fetal development [3].

Exposure to Particulate Matter (PM) can affect prenatal maternal health through a variety of mechanisms, including increased Reactive Oxygen Species (ROS), which stimulates macrophages to secrete cytokines TNF- α and NF κ B which will further accelerate the process of apoptosis in placenta [2].

Currently, interest in stem cells for the treatment of various diseases is increasing. The MarrowStem Bone Cell is utilized for therapy because it has a self-renewing properties, so that therapy need not be repeated. In an teratogenic case, Bone Marrow Stem Cell administration can reduce the inflammation, apoptosis and congenital abnormalities [4].

Therefore, the utilization of Bone Marrow Stem Cell for therapy in the teratogenic case requires a biomolecular reproduction study to reveal the mechanisms of disorders in placentation during pregnancy as a basis of proving the study scientifically. This study used rats as an animal model of teratogenic particulate matter merely for ethical consideration and smooth execution of the study.

MATERIALS AND METHODS

This study used 2 groups. The control group (K) had 15 pregnant rats exposed to carbon black powder with a dose of 532 mg / m³ on per-inhalation of 4 hours / day during pregnancy of 7-18 days without Rat Bone Marrow Stem Cell therapy. The treatment group (P) had 15 pregnant rats exposed to carbon black powder with a dose of 532 mg / m³ on a per-inhalation of 4 hours / day during pregnancy of 7-18 days with Rat Bone Marrow Stem Cell therapy with a dosage of 106 / head. The intravenous injections were made on the tails. On day 19 the termination and placental tissue examination to see NF κ B expression by immunohistochemistry were done.

RESULTS AND DISCUSSION


Each sample was semi-qualitatively assessed using modified Remmele method. The assessment result of the expression of NF κ B in placenta using modified method of semi-qualitative Remmele Index Scale (IRS) then tested using Kruskal Wallis was observably very different, then continued with Mann-Whitney test for the two treatments which can be seen in table 1.

Table 1 shows that there is a significant result on the treatment group (exposed to Carbon Black and given Stem-Cell therapy) compared to the control group (exposed to Carbon Black). It shows that Rat Bone Marrow Stem Cell can decrease placental NF- κ B expression [5].

Table 1. Score of placental Nf- κ B expression

Immunohistochemistry	Group	Comparing Group	Significance
Nf- κ B	Control	Treatment	0.001**

**Significance $p < 0.01$



Picture 1. Picture A dan B show Nf κ B expression of placental tropoblast cells. Control group (picture A) shows negative expression (arrow). Treatment group (picture B) shows possitive expression shown by brownish color. (Coloring Imunohisto-chemistry,1000x)

CONCLUSION

The result of rat bone marrow stem cell therapy administration on pregnant white rats exposed to particulate matter showed a decrease in placental NF- κ B expression.

REFERENCES

- [1] Balai Besar Teknik Kesehatan Lingkungan dan Pemberantasan Penyakit Menular/BBTKL dan P2M. 2009. Laporan: Situasi dan kecenderungan parameter pencemar udara dan air minum serta potensi resiko gangguan kesehatan di Jawa Timur 2006-2008. Surabaya.
- [2] Biri A, Bozkurt N, Turp A, Kavutcu M, Himmetoglu O, Durak I. 2007. Role of oxidative stress in intrauterine growth restriction. *Ginecol Obstet Invest* 64(4): 187-192.
- [3] Dejmek J, Selevan SG, Benes I, Solansky I, Sram RJ. 1999. Fetal growth and maternal exposure to particulate matter during pregnancy. *Environmental Health Perspectives* 107 960; 475-480.
- [4] Lee, H.J., K. Selesmiami, Y. Niikura, T. Niikura, R. Klein D.M. Dombkowski and J.L. Tilly. 2007. Bone Marrow Transplantation Generates Immature Oocytes and Rescues Ling Term Fertility in a Preclinical Mouse Model of Chemotherapy Induced Premature Ovarian Failure. 25(22):3198-3204
- [5] Wang, Yun-Ying, K.Sui, H. Li., and H. Ma. 2009. The Effects of Lead Exposure on Placental NF κ B Expression and The Consequences for Gestation. *J. Reprotox* 27:190-195

Effectiveness of Rat Bone Marrow Stem Cell Therapy to Rattus novergicus by Teratogenic Model of Particulate Matter on Expression of Kappa Beta (NFk(I) Nuclear Factor on Placenta

ORIGINALITY REPORT

25%

SIMILARITY INDEX

18%

INTERNET SOURCES

13%

PUBLICATIONS

19%

STUDENT PAPERS

PRIMARY SOURCES

1

jurnal.unsyiah.ac.id

Internet Source

12%

2

repository.unair.ac.id

Internet Source

4%

3

Submitted to iGroup

Student Paper

4%

4

Geir Bjørklund, Maryam Dadar, Joachim Mutter, Jan Aaseth. "The toxicology of mercury: Current research and emerging trends", Environmental Research, 2017

Publication

2%

5

www.ncbi.nlm.nih.gov

Internet Source

1%

6

Submitted to Universitas Airlangga

Student Paper

1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Effectiveness of Rat Bone Marrow Stem Cell Therapy to Rattus novergicus by Teratogenic Model of Particulate Matter on Expression of Kappa Beta (NFk(I) Nuclear Factor on Placenta

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2
